

# Hardy-Littlewood-Stein-Weiss inequality in the generalized Morrey spaces

Aygun N. Huseynova

Azerbaijan State Oil and Industry University

Baku, Azerbaijan

Email:hasanova\_a@inbox.ru

(Received: February 18, 2021, Accepted: June 29, 2021)

## Abstract

We consider generalized weighted Morrey spaces  $\mathcal{M}^{p,\omega,|\cdot|^\gamma}(\mathbb{R}^n)$  and a general function  $\omega(x, r)$  defining the Morrey-type norm. We prove the Riesz potential  $I_\alpha$  is bounded from the weighted Morrey space  $\mathcal{M}^{p,\omega_1,|\cdot|^\gamma}(\mathbb{R}^n)$  to  $\mathcal{M}^{q,\omega_2,|\cdot|^\mu}(\mathbb{R}^n)$ ,  $1 < p < \frac{n}{\alpha}$ ,  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$ ,  $\alpha p - n < \gamma < n(p-1)$ ,  $\mu = \frac{q\gamma}{p}$ .

## 1 Introduction

Morrey spaces were introduced by C.B. Morrey in 1938 in connection with certain problems in elliptic partial differential equations and calculus of variations (see [21]), they are defined by the norm

$$\|f\|_{\mathcal{L}^{p,\lambda}} := \sup_{x,r>0} r^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x,r))},$$

where  $0 \leq \lambda < n$ ,  $1 \leq p < \infty$ . In the theory of partial differential equations, together with weighted Lebesgue spaces, Morrey spaces  $\mathcal{L}^{p,\lambda}(\Omega)$  play an important

---

**Keywords and phrases:** Maximal operator, fraction maximal operator, Riesz potential, generalized Morrey space, Hardy-Littlewood-Stein-Weiss type estimate, BMO space

**2020 AMS Subject Classification:** Primary 42B20, 42B25, 42B35

role. Later, Morrey spaces found important applications to Navier-Stokes ([20], [32]) and Schrodinger ([24]), [26], [27], [30], [31]) equations, elliptic problems with dicontinuous coefficients ([4], [6]) and potential theory ([1], [2]). An exposition of the Morrey spaces can be found in the books [7] and [18].

Generalized Morrey spaces of such a kind in the case of contant  $p$  were studied in [3], [19], [22], [23]. In [10] there was proved the boundedness of the maximal operator, singular integral operator and the potential operators in generalized variable exponent Morrey spaces.

The results on the boundedness of potential operators and classical Calderon-Zygmund singular operators go back to [1] and [25], respectively, while the boundeness of the maximal operator in the Euclidean setting was proved in [5].

Hardy-Littlewood-Stein-Weiss inequality in the Lebesgue spaces was proved in H. G. Hardy and J. E. Littlewood [16] in the one-dimensional case and to E. M. Stein and G. Weiss [29] in the case  $n > 1$ .

One of the most important variants of the Hardy-Littlewood maximal function is the so-called fractional maximal function defined by the formula

$$M_\alpha f(x) = \sup_{t>0} |B(x, t)|^{-1+\alpha/n} \int_{B(x, t)} |f(y)| dy, \quad 0 \leq \alpha < n,$$

where  $|B(x, t)|$  is the Lebesgue measure of the ball  $B(x, t)$ .

It coincides with the Hardy-Littlewood maximal function  $Mf \equiv M_0 f$  and is intimately related to the Riesz potential

$$I_\alpha f(x) = \int_{\mathbb{R}^n} \frac{f(y) dy}{|x - y|^{n-\alpha}}, \quad 0 < \alpha < n.$$

The paper is organized as follows. In Section 2 we provide necessary preliminaries on variable exponent Lebesgue and Morrey spaces. In Section 3 we treat Riesz potentials.

The main results are given in Theorems 3.1, 3.2, 3.3, 3.4.

## 2 Preliminaries on Lebesgue and Morrey spaces

$L_p \equiv L_p(\mathbb{R}^n)$  is the space of all classes of measurable functions  $f$  with finite norm

$$\|f\|_{L_p} = \left( \int_{\mathbb{R}^n} |f(x)|^p dx \right)^{\frac{1}{p}}, \quad 1 \leq p < \infty$$

and also  $\omega L_p(\mathbb{R}^n)$ , the weak  $L_p$  space defined as the set of all measurable function  $f$  on  $\mathbb{R}^n$  with the following finite norms

$$\|f\|_{\omega L_p} = \sup_{r>0} r |\{x \in \mathbb{R}^n : |f(x)| > r\}|^{1/p}, \quad 1 \leq p < \infty.$$

For  $p = \infty$  the space  $L_\infty(\mathbb{R}^n)$  is defined by means of the usual modification

$$\|f\|_{L_\infty} = \operatorname{ess\,sup}_{x \in \mathbb{R}^n} |f(x)|.$$

Let  $L_{p,\omega}(\mathbb{R}^n)$  be the space of measurable functions on  $\mathbb{R}^n$  with finite norm

$$\|f\|_{L_{p,\omega}} = \|f\|_{L_{p,\omega}(\mathbb{R}^n)} = \left( \int_{\mathbb{R}^n} |f(x)|^p \omega(x) dx \right)^{1/p}, \quad 1 \leq p < \infty.$$

and for  $p = \infty$  the space  $L_{\infty,\omega}(\mathbb{R}^n) = L_\infty(\mathbb{R}^n)$ ,

**Definition 2.1.** *The weight function  $\omega$  belongs to the class  $A_p(\mathbb{R}^n)$ , for  $1 \leq p < \infty$ , if the following statement*

$$\sup_{x \in \mathbb{R}^n, r>0} \frac{1}{|B(x,r)|} \int_{B(x,r)} \omega(y) dy \left( \frac{1}{|B(x,r)|} \int_{B(x,r)} \omega^{-\frac{1}{p-1}}(y) dy \right)^{p-1}$$

is finite and  $\omega$  belongs to  $A_1(\mathbb{R}^n)$ , if there exists a positive constant  $C$  such that for any  $x \in \mathbb{R}^n$  and  $r > 0$

$$|B(x,r)|^{-1} \int_{B(x,r)} \omega(y) dy \leq C \operatorname{ess\,sup}_{y \in B(x,r)} \frac{1}{\omega(y)}.$$

The following theorem was proved in [29].

**Theorem 2.1.** *Let  $0 < \alpha < n$ ,  $1 < p < \frac{n}{\alpha}$ ,  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$ ,  $\alpha p - n < \gamma < n(p-1)$ ,  $\mu = \frac{q\gamma}{p}$ . Then the operator  $I^\alpha$  is bounded from  $L_{p,|\cdot|^\gamma}(\mathbb{R}^n)$  to  $L_{q,|\cdot|^\mu}(\mathbb{R}^n)$ .*

Let  $M^\#$  be the sharp maximal function defined by

$$M^\# f(x) = \sup_{r>0} |B(x,r)|^{-1} \int_{B(x,r)} |f(y) - f_{B(x,r)}| dy.$$

where  $f_{B(x,t)}(x) = |B(x,t)|^{-1} \int_{B(x,t)} f(y) dy$ .

**Definition 2.2.** We define the  $BMO(\mathbb{R}^n)$  space as the set of all locally integrable functions  $f$  with finite norm

$$\|f\|_{BMO} = \sup_{x \in \mathbb{R}^n, r > 0} |B(x, r)|^{-1} \int_{B(x, r)} |f(y) - f_{B(x, r)}| dy$$

or

$$\|f\|_{BMO} = \inf_C \sup_{x \in \mathbb{R}^n, r > 0} |B(x, r)|^{-1} \int_{B(x, r)} |f(y) - C| dy.$$

where  $f_{B(x, t)}(x) = |B(x, t)|^{-1} \int_{B(x, t)} f(y) dy$ .

**Definition 2.3.** We define the  $BMO_{p, \omega}(\mathbb{R}^n)$  ( $1 \leq p < \infty$ ) space as the set of all locally integrable functions  $f$  with finite norm

$$\|f\|_{BMO_{p, \omega}} = \sup_{x \in \mathbb{R}^n, r > 0} \frac{\|(f(\cdot) - f_{B(x, r)})\chi_B(x, r)\|_{L_{p, \omega}(\mathbb{R}^n)}}{\|\chi_B(x, r)\|_{L_{p, \omega}(\mathbb{R}^n)}}.$$

**Theorem 2.2** (17, Theorem 4.4). Let  $1 \leq p < \infty$  and  $\omega$  be a Lebesgue measurable function. If  $\omega \in A_p(\mathbb{R}^n)$ , then the norms  $\|\cdot\|_{BMO_{p, \omega}}$  and  $\|\cdot\|_{BMO}$  are mutually equivalent.

We find it convenient to define the generalized Morrey spaces in the form as follows

**Definition 2.4.** Let  $1 \leq p < \infty$ . The generalized Morrey space  $M^{p, \omega}(\mathbb{R}^n)$  and generalized weighted Morrey space  $M^{p, \omega, |\cdot|}(\mathbb{R}^n)$  are defined by the norms

$$\|f\|_{M^{p, \omega}} = \sup_{x \in \mathbb{R}^n, r > 0} \frac{r^{-\frac{n}{p}}}{\omega(x, r)} \|f\|_{L_p(B(x, r))},$$

$$\|f\|_{M^{p, \omega, |\cdot|} \gamma} = \sup_{x \in \mathbb{R}^n, r > 0} \frac{r^{-\frac{n}{p}}}{\omega(x, r)} \|f\|_{L_{p, |\cdot|} \gamma(B(x, r))},$$

Everywhere in the sequel we assume that

$$\inf_{x \in \mathbb{R}^n, r > 0} \omega(x, r) > 0$$

which makes the space  $M^{p, \omega} \mathbb{R}^n$  nontrivial. Note that when  $p$  is constant, in the case of  $\omega(x, r) \equiv \text{const} > 0$ , we have the space  $L^\infty(\mathbb{R}^n)$ .

### 3 Riesz potential operator in the spaces $M^{p,\omega,|\cdot|^\gamma}(\mathbb{R}^n)$

**Theorem 3.1.** *Let  $0 < \alpha < n$ ,  $1 < p < \frac{n}{\alpha}$ ,  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$ ,  $\alpha p - n < \gamma < n(p-1)$ ,  $\mu = \frac{\alpha\gamma}{p}$ . Then*

$$\|I^\alpha f\|_{L_{q,|\cdot|^\mu}(B(x,t))} \leq C t^{\frac{n}{q} + \frac{\gamma}{p}} \int_t^\infty s^{-\frac{n}{q} - \frac{\gamma}{p} - 1} \|f\|_{L_{p,|\cdot|^\gamma}(B(x,s))} ds, \quad t > 0 \quad (3.1)$$

where  $C$  does not depend on  $f, x$  and  $t$ ,

*Proof.* We represent  $f$  as

$$f = f_1 + f_2, \quad f_1(y) = f(y)\chi_{B(x,2t)}(y), \quad f_2(y) = f(y)\chi_{\mathbb{R}^n \setminus B(x,2t)}(y), \quad t > 0, \quad (3.2)$$

and have

$$I^\alpha f(x) = I^\alpha f_1(x) + I^\alpha f_2(x)$$

By Theorem 2.1 we obtain

$$\|I^\alpha f_1\|_{L_{q,|\cdot|^\mu}(B(x,t))} \leq \|I^\alpha f_1\|_{L_{q,|\cdot|^\mu}(\mathbb{R}^n)} \leq C \|f_1\|_{L_{p,|\cdot|^\gamma}(\mathbb{R}^n)} = C \|f_1\|_{L_{p,|\cdot|^\gamma}(B(x,2t))}.$$

Then

$$\|I^\alpha f_1\|_{L_{q,|\cdot|^\mu}(B(x,t))} \leq C \|f\|_{L_{p,|\cdot|^\gamma}(B(x,2t))},$$

where the constant  $C$  is independent of  $f$ .

Taking into account that

$$\|f\|_{L_{p,|\cdot|^\gamma}(B(x,2t))} \leq C t^{\frac{n}{q} + \frac{\gamma}{p}} \int_t^\infty s^{-\frac{n}{q} - \frac{\gamma}{p} - 1} \|f\|_{L_{p,|\cdot|^\gamma}(B(x,s))} ds$$

we get

$$\|I^\alpha f_1\|_{L_{q,|\cdot|^\mu}(B(x,t))} \leq C t^{\frac{n}{q} + \frac{\gamma}{p}} \int_t^\infty s^{-\frac{n}{q} - \frac{\gamma}{p} - 1} \|f\|_{L_{p,|\cdot|^\gamma}(B(x,s))} ds \quad (3.3)$$

When  $|x - z| \leq t$ ,  $|z - y| \geq 2t$ , we have  $\frac{1}{2}|z - y| \leq |x - y| \leq \frac{3}{2}|z - y|$ , and therefore

$$|I^\alpha f_2(x)| \leq \int_{\mathbb{R}^n \setminus B(x,2t)} |z - y|^{\alpha-n} |f(y)| dy \leq C \int_{\mathbb{R}^n \setminus B(x,2t)} |x - y|^{\alpha-n} |f(y)| dy.$$

We choose  $\beta > \frac{n}{q}$  and obtain

$$\begin{aligned}
& \int_{\mathbb{R}^n \setminus B(x, 2t)} |x - y|^{\alpha-n} |f(y)| dy \\
&= \beta \int_{\mathbb{R}^n \setminus B(x, 2t)} |x - y|^{\alpha-n+\beta} |f(y)| dy \left( \int_{|x-y|}^{\infty} s^{-\beta-1} ds \right) dy \\
&= \beta \int_2^{\infty} t S^{-\beta-1} \left( \int_{\{y \in \mathbb{R}^n : 2t \leq |x-y| \leq s\}} |x - y|^{\alpha-n+\beta} |f(y)| dy \right) ds \\
&\leq C \int_2^{\infty} t S^{-\beta-1} \|f\|_{L_{q,|\cdot|}^{\gamma}(B(x,s))} \| |x - y|^{\alpha-n+\beta} \|_{L_{p'(\cdot),|\cdot|^{\gamma/(1-p)}}(B(x,s))} ds \\
&\leq C \int_2^{\infty} t S^{\alpha - \frac{n}{q} - \frac{\gamma}{p} - 1} \|f\|_{L_{p,|\cdot|^{\gamma}}(B(x,s))} ds
\end{aligned}$$

Hence

$$\|I^{\alpha} f_2\|_{L_{q,|\cdot|^{\mu}}(B(x,t))} \leq C \int_2^{\infty} t s^{\frac{n}{q} - \frac{\gamma}{p} - 1} \|f\|_{L_{p,|\cdot|^{\gamma}}(B(x,s))} ds \|X_{B(x,t)}\|_{L_{q,|\cdot|^{\mu}}(\mathbb{R}^n)}$$

Therefore

$$\|I^{\alpha} f_2\|_{L_{q,|\cdot|^{\mu}}(B(x,t))} \leq C t^{\frac{n}{q} - \frac{\gamma}{p}} \int_2^{\infty} 2ts^{-\frac{n}{q} - \frac{\gamma}{p} - 1} \|f\|_{L_{p,|\cdot|^{\gamma}}(B(x,s))} ds \quad (3.4)$$

which together (3.3) yields (3.1).  $\square$

**Theorem 3.2.** Let  $0 < \alpha < n$ ,  $1 < p < \frac{n}{\alpha}$ ,  $-\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$ ,  $\alpha p - n < \gamma < n(p-1)$ ,  $\mu = \frac{\alpha\gamma}{p}$  and the function  $\omega_1(x, r)$  and  $\omega_2(x, r)$  fulfill the condition

$$\int_r^{\infty} t^{\alpha - \frac{\gamma}{p}} \omega_1(x, t) \frac{dt}{t} \leq C r^{-\frac{\gamma}{p}} \omega_2(x, r).$$

Then the operators  $M^{\alpha}$  and  $I^{\alpha}$  are bounded from  $M^{p, \omega_1(\cdot), |\cdot|^{\gamma}}(\mathbb{R}^n)$  to  $M^{q, \omega_2(\cdot), |\cdot|^{\mu}}(\mathbb{R}^n)$ .

*Proof.* Let  $f \in M^{p, \omega_1(\cdot), |\cdot|^{\gamma}}(\mathbb{R}^n)$ . As usual, when estimating the norm

$$\|I^{\alpha} f\|_{M^{q, \omega_2(\cdot), |\cdot|^{\mu}}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n, t > 0} \frac{t^{-\frac{n}{q}}}{\omega_2(x, t)} \|I^{\alpha} \int_{XB(x,t)} \|L_{q,|\cdot|^{\mu}}(\mathbb{R}^n)\|. \quad (3.6)$$

We estimate  $\|I^\alpha f XB(x, t)\|_{L_{q,|\cdot|}^\mu(\mathbb{R}^n)}$  in (3.6) by means of Theorem 3.1 and obtain

$$\begin{aligned} & \|I^\alpha f\|_{M^{q,\omega_2,|\cdot|}^\mu(\mathbb{R}^n)} \\ & \leq C \sup_{x \in \mathbb{R}^n, t > 0} \frac{t^{-\frac{\gamma}{p}}}{\omega_2(x, t)} \int_t^\infty r^{\frac{n}{q} - \frac{\gamma}{p} - 1} \|f\|_{L_{p,|\cdot|,\gamma}(B(x, r))} dr \\ & \leq C \|f\|_{M^{p,\omega_1,|\cdot|}^\gamma(\mathbb{R}^n)} \sup_{x \in \mathbb{R}^n, t > 0} \frac{t^{\frac{\gamma}{p}}}{\omega_2(x, t)} \int_t^\infty \frac{r^{\alpha - \frac{\gamma}{p}} \omega_1(x, r)}{r} dr. \end{aligned}$$

It remains to make use of condition (3.5).  $\square$

**Theorem 3.3.** Let  $0 < \alpha < n$ ,  $1 < p < \frac{n}{\alpha}$ ,  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$ ,  $\gamma, \mu$  satisfy condition

$$0 \leq \gamma < \frac{n}{p'}, \quad \mu = \frac{\gamma}{p}, \quad (3.7)$$

and let  $\omega(x, t)$  satisfy condition

$$r^\alpha \omega(x, r) \leq C, \quad (3.8)$$

Then the operators  $M^\alpha$  is bounded from  $M^{p,\omega(\cdot),|\cdot|}^\gamma(\mathbb{R}^n)$  to  $L_{\infty,|\cdot|}^\mu(\mathbb{R}^n)$ .

*Proof.* Let  $x \in \mathbb{R}^n$  and  $r > 0$ . By the Holder inequality we get successively

$$\begin{aligned} & r^{\alpha-n} \int_{B(x,r)} |f(y)| dy \\ & \leq C r^{\alpha-n} r^{\frac{n}{p}} \omega(x, r) r^{-\frac{n}{p}} \omega^{-1}(x, r) \|f\|_{L_{p,|\cdot|,\gamma}(B(x, r))} \|XB(x, r)\|_{L_{p',|\cdot|}^{\gamma/(1-p)}} \\ & \leq C r^\alpha \omega(x, r) |x|^{-\frac{\gamma}{p}} \|f\|_{M^{p,\omega(\cdot),|\cdot|}^\gamma} \leq C |x|^{-\frac{\gamma}{p}} \|f\|_{M^{p,\omega(\cdot),|\cdot|}^\gamma}. \end{aligned}$$

$\square$

**Theorem 3.4.** Let  $0 < \alpha < n$ ,  $1 < p < \frac{n}{\alpha}$ ,  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$ ,  $\alpha p - n < \gamma < n(p-1)$ ,  $\gamma, \mu$  satisfy condition (3.7),  $0 < \alpha < n$  and let  $\omega(x, t)$  satisfy condition (3.8).

Then the operators  $I^\alpha$  is bounded from  $M^{p,\omega(\cdot),|\cdot|}^\gamma(\mathbb{R}^n)$  to  $BMO(\mathbb{R}^n)$ ,

*Proof.* Let  $f \in M^{p,\omega(\cdot),|\cdot|^\gamma}(\mathbb{R}^n)$ . In [1] was proved

$$M^t(I^\alpha f)(x) \leq CM^\alpha f(x). \quad (3.9)$$

The proof Theorem 3.4, by the Theorem 3.3 and inequality (3.9).  $\square$

## References

- [1] D. R. Adams, *A note on Riesz potentials*, Duke Math., **42**(1975), 765-778.
- [2] D. R. Adams, *Choquet integrals in potential theory*, Publ. Mat., **42**(1998), 3-66.
- [3] V. I. Burenkov and H. V. Guliyev, *Necessary and sufficient conditions for boundedness of the maximal operator in the local Morrey-type spaces*, Studia Mathematica, **163**(2) (2004), 157-176.
- [4] L. Caffarelli, *Elliptic second order equations*, Rend. Sem. Mat. Fis. Milano, **58** (1990), 253-284.
- [5] F. Chiarenza and M. Frasca, *Morrey spaces and Hardy-Littlewood maximal function*, Rend. Math., **7** (1987), 273-279.
- [6] G. Di Fazio, D. K. Palagachev and M. A. Ragusa, *Global Morrey regularity of strong solutions to the Dirichlet problem for elliptic equations with discontinuous coefficients*, J. Funct. Anal., **166** (1999), 179-196.
- [7] M. Giaquinta, *Multiple integrals in the calculus of variations and nonlinear elliptic systems*, Princeton Univ. Press, Princeton, NJ, 1983.
- [8] V. S. Guliyev, *Integral operators on function spaces on the homogeneous groups and on domains in  $\mathbb{R}^n$* , Doctor's degree dissertation, Moscow, Mat. Inst. Steklov, (1994), 1-329 (Russian).
- [9] V. S. Guliyev, *Function spaces, integral operators and two weighted inequalities on homogeneous groups*, Some applications, Baku, (1999), 1-332 (Russian).
- [10] V. S. Guliyev, J. J. Hasanov and S. G. Samko, *Boundedness of the maximal, potential and singular operators in the generalized variable exponent Morrey spaces*, Mathematica Scandinavica, **107**, (2010), 285-304.



- [11] V. S. Guliyev and R. Ch Mustafayev, *Integral operators of potential type in spaces of homogeneous type*, (Russian) Doklady Ross, Akad. Nauk., **354**(6), (1997), 730-732.
- [12] V. S. Guliyev and R. Ch Mustafayev, *Fractional integrals in spaces of functions defined on spaces of homogeneous type*, (Russian) Anal. Math., **24**(3), (1998), 181-200.
- [13] A. I. Guseinov and Kh. Sh. Mukhtarov, *Introduction to the theory of nonlinear singular integral equations*, (Russian) Moscow, Nauka, 1980.
- [14] J. J. Hasanov, *Hardy-Littlewood-Stein-Weiss inequality in the variable exponent Morrey spaces*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb., **39**, (2013), 47-62.
- [15] J. J. Hasanov and A. M. Musayev, *Oscillatory integral operators and their commutators in modified weighted Morrey spaces with variable exponent*, International Journal of Applied Mathematics, **32**(3), (2019), 521-535.
- [16] H.G. Hardy and J.E. Littlewood, *Some properties of fractional integrals*, I, Math. Z., **27**(4) (1928), 565-606.
- [17] K-P. Ho, *Singular integral operators, John-Nirenberg inequalities and Tribel-Lizorkin type spaces on weighted Lebesgue spaces with variable exponents*, Revista De La Union Matematica Argentina, **57**(1) (2016), 85-101.
- [18] A. Kufner, O. John and S. Fucik, *Function spaces*, Noordhoff International Publishing Leyden, Publishing HOUse Czechoslovak Academy of Sciences, Prague, (1977).
- [19] K. Kurata, S. Nishigaki and S. Sugano, *Boundedness of Integral Operators on Generalized Morrey Spaces and Its Application to Schrodinger operators*, Proc. AMS., **128**(4) (1999), 1125-1134 .
- [20] A. L. Mazzucato, *Besov-Morrey spaces: function space theory and applications to non-linear PDE*, Trans. Amer. Math. Soc., **355**, (2003), 1297-1364.
- [21] C. B. Morrey, *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc., **43** (1938), 126-166.
- [22] T. Mizuhara, *Boundedness of some classical operators on generalized Morrey spaces*, Harmonic Analysis (S. Igari, Editor), ICM 90 Satellite Proceedings, Springer - Verlag, Tokyo, (1991), 183-189.

- [23] E. Nakai, *Hardy-Littlewood maximal operators, singular integral operators and Riesz potential on generalized Morrey spaces*, Math. Nachr, **166**, (1994), 95-103.
- [24] C. Perez, *Two weighted norm inequalities for Riesz potentials and uniform  $L_p$ -weighted Sobolev inequalities*, Indiana Univ. Math. J., **39** (1990) 3144.
- [25] J. Peetre, *On the theory of  $\mathcal{L}^{p,\lambda}$  spaces*, J. Funct. Anal., **4**, (1969), 71-87.
- [26] A. Ruiz and L. Vega, *Unique continuation for Schrodinger operators with potential in Morrey spaces*, Publ. Mat., **35**, (1991), 291-298.
- [27] A. Ruiz and L. Vega, *On local regularity of Schrodinger equations*, Int. Math. Res. Notices, **1993**(1), (1993), 13-27.
- [28] E. M. Stein, *Harmonic Analysis: Real-variable Methods, Orthogonality and Oscillatory integrals*, Princeton Univ. Press, Princeton, N.J., (1993).
- [29] E. M. Stein and G. Weiss, *Fractional integrals on  $n$ -dimensional Euclidean space*, J. Math. and Mech., **7**(4), (1958), 503-514.
- [30] Z. Shen, *The periodic Schrodinger operators with potentials in the Morrey class*, J. Funct. Anal., **193**, (2002), 314-345.
- [31] Z. Shen, *Boundary value problems in Morrey spaces for elliptic systems on Lipschitz domains*, Amer. J. Math., **125**, (2003), 1079-1115.
- [32] M. E. Taylor, *Analysis on Morrey spaces and applications to Navier-Stokes and other evolution equations*, Comm. Partial Differential Equations, **17**, (1992), 1407-1456.